arXiv:1302.4105v1 [hep-th] 17 Feb 2013

SLAC-PUB-15366

Threefold Complementary Approach to Holographic QCD

Stanley J. Brodsky,[1, *] Guy F. de Téramond,[2, †] and Hans Günter Dosch[3, ‡]

[1] *SLAC National Accelerator Laboratory,*
Stanford University, Stanford, California 94309, USA
[2] *Universidad de Costa Rica, San José, Costa Rica*
[3] *Institut für Theoretische Physik, Philosophenweg 16, D-6900 Heidelberg, Germany*

(Dated: February 19, 2013)

Abstract

A complementary approach, based on higher dimensional anti-de Sitter (AdS) space, light-front quantization, and the invariance properties of the one-dimensional representation of the full conformal group, provides a nonperturbative model which successfully incorporates salient features of hadronic physics, including confinement, linear Regge trajectories, and results which are conventionally attributed to spontaneous chiral symmetry breaking.

A remarkable holographic correspondence between the equations of motion in anti–de Sitter (AdS) space and the Hamiltonian equations of motion for relativistic light hadron bound-states in physical space-time was derived in Ref. [1]. To a first semiclassical approximation, light-front QCD is formally equivalent to the equations of motion on a fixed gravitational background asymptotic to AdS_5. The confinement properties are encoded in a dilaton $\varphi(z)$ which introduces a length scale in the AdS action and leads to conformal symmetry breaking. In this paper we shall show that similar results follow from an elegant construction of Hamiltonian operators by V. de Alfaro, S. Fubini and G. Furlan (dAFF) [2], based on the invariance properties of a field theory in one dimension under the full conformal group. In contrast with the holographic approach, the construction of dAFF retains conformal invariance of the action despite the presence of a fundamental length scale. In addition, we will use complementary arguments from AdS space, light-front (LF) quantization, and the dAFF construction to show that the form of the dilaton profile in AdS space $\varphi(z) \propto z^s$ is constrained to have the specific power $s = 2$. This profile leads not only to linear Regge trajectories [3], but it also leads uniquely to a massless pion in the chiral limit [4].

The central problem for solving QCD using the LF Hamiltonian method can be reduced to the derivation of the effective interaction in the LF wave equation which acts on the valence sector of the theory and has, by definition, the same eigenvalue spectrum as the initial Hamiltonian problem. In order to carry out this program one must systematically express the higher Fock components as functionals of the lower ones. The method has the advantage that the Fock space is not truncated, and the symmetries of the Lagrangian are preserved [5].

In the limit of zero quark masses the longitudinal modes decouple from the invariant LF Hamiltonian equation $H_{LF}|\phi\rangle = M^2|\phi\rangle$ with $H_{LF} = P_\mu P^\mu = P^- P^+ - \vec{P}_\perp^2$. The generators $P = (P^-, P^+, \vec{P}_\perp)$, $P^\pm = P^0 \pm P^3$, are constructed canonically from the QCD Lagrangian by quantizing the system on the light-front at fixed LF time x^+, $x^\pm = x^0 \pm x^3$ [6]. The LF Hamiltonian P^- generates the LF time evolution

$$P^-|\phi\rangle = i\frac{\partial}{\partial x^+}|\phi\rangle, \tag{1}$$

whereas the LF longitudinal P^+ and transverse momentum $\vec{P}_\perp$ are kinematical generators. We obtain the wave equation [1]

$$\left(-\frac{d^2}{d\zeta^2} - \frac{1-4L^2}{4\zeta^2} + U\left(\zeta, J\right)\right)\phi_{n,J,L} = M^2\phi_{n,J,L}, \tag{2}$$

a relativistic single-variable LF Schrödinger equation. This equation describes the spectrum of mesons as a function of n, the number of nodes in ζ, the total angular momentum J, which represent the maximum value of $|J^z|$, $J = \max|J^z|$, and the internal orbital angular momentum of the constituents $L = \max|L^z|$. The variable z of AdS space is identified with the LF boost-invariant transverse-impact variable ζ [7], thus giving the holographic variable a precise definition in LF QCD [1, 7]. For a two-parton bound state $\zeta^2 = x(1-x)b_\perp^2$, where x is the longitudinal momentum fraction and $b_\perp$ is the transverse-impact distance between the quark and antiquark. In the exact QCD theory U is related to the two-particle irreducible $q\bar{q}$ Green's function.

Recently we have derived wave equations for hadrons with arbitrary spin starting from an effective action in AdS space [8]. An essential element is the mapping of the higher-dimensional equations to the LF Hamiltonian equation found in Ref. [1]. This procedure allows a clear distinction between the kinematical and dynamical aspects of the LF holographic approach to hadron physics. Accordingly, the non-trivial geometry of pure AdS space encodes the kinematics, and the additional deformations of AdS encode the dynamics, including confinement [8].

For $d = 4$ one finds the LF potential [9]

$$U(\zeta, J) = \frac{1}{2}\varphi''(\zeta) + \frac{1}{4}\varphi'(\zeta)^2 + \frac{2J-3}{2\zeta}\varphi'(\zeta), \tag{3}$$

provided that the product of the AdS mass m and the AdS curvature radius R are related to the total and orbital angular momentum, J and L respectively, according to $(mR)^2 = -(2-J)^2 + L^2$. The critical value $J = L = 0$ corresponds to the lowest possible stable solution, the ground state of the LF Hamiltonian in agreement with the AdS stability bound $(mR)^2 \geq -4$ [10].

The choice $\varphi(z) = \lambda z^2$ leads through (3) to the LF potential $U(\zeta, J) = \lambda^2\zeta^2 + 2\lambda(J-1)$, and Eq. (2) yields the mass spectrum for the mesons

$$M_{n,J,L}^2 = 2\lambda\left(2n + J + L\right). \tag{4}$$

This result not only implies linear Regge trajectories, but also a massless pion as well as the relation between the ρ and a_1 mass usually obtained from the Weinberg sum rules [11]

$$m_\pi^2 = M_{0,0,0}^2 = 0,\ m_\rho^2 = M_{0,1,0}^2 = 2\lambda,\ m_{a_1}^2 = M_{0,1,1}^2 = 4\lambda. \tag{5}$$

If one starts with a dilaton profile of the general form

$$\varphi_s(z) = \lambda^{s/2}\, z^s, \tag{6}$$

for $\lambda > 0$ and arbitrary values of s, the existence of a massless pion in the limit of massless quarks determines uniquely the value $s = 2$. To show this, we use the stationarity of bound-state energies with respect to variation of parameters [12]. For $J = 0$ the LF potential (3) for the dilaton profile (6) is

$$U_s(\zeta) = \frac{1}{2} s(s-4)\lambda^{s/2}\zeta^{s-2} + \frac{1}{4} s^2 \lambda^s\, \zeta^{2s-2}. \tag{7}$$

We make a variational ansatz with the test function $\tilde{\phi}(\zeta,\alpha) = \sqrt{2\lambda\alpha\,\zeta}\; e^{-\lambda\alpha\,\zeta^2/2}$ where $\langle\tilde{\phi}(\zeta,\alpha)|\tilde{\phi}(\zeta,\alpha)\rangle = 1$. The minimum of the expectation value of the LF Hamiltonian $\langle M_s^2(\alpha)\rangle = \langle\tilde{\phi}(\zeta,\alpha)|H_{LF}|\tilde{\phi}(\zeta,\alpha)\rangle$

$$\begin{aligned}\langle M_s^2(\alpha)\rangle &= \int d\zeta\, \tilde{\phi}(\zeta,\alpha)\left(-\frac{d^2}{d\zeta} - \frac{1}{4\zeta^2} + U_s(\zeta)\right)\tilde{\phi}(\zeta,\alpha) \\ &= \lambda\alpha\left(1 + (s-4)\,\Gamma(1+s/2)\,\alpha^{-s/2} + \frac{s}{4}\,\Gamma(1+s)\,\alpha^{-s}\right),\end{aligned} \tag{8}$$

with respect to the parameter α, $\frac{\partial}{\partial\alpha}\langle M_s^2(\alpha)\rangle = 0$, gives an upper bound for the lowest eigenvalue $n = J = L = 0$ for a given value of s.



FIG. 1: Stationary value of $\langle M_s^2(\alpha)\rangle/\lambda$ as a function of the power s in the dilaton profile (6). The circles are the results of numerical solution for $s > 2$. The numerical errors are smaller than the size of the circles in the figure.

In Fig. 1 we display the stationary value $M_s^2/\lambda = \min_\alpha\langle M_s^2(\alpha)\rangle/\lambda$ as a function of s. Since for $s < 2$ the upper bound is negative, the exact value of the ground state mass M_{000}^2

has also to be negative, thus not a possible physical state. For $s = 2$ the stationary value M_s^2 is the exact eigenvalue and has the value zero. For $s > 2$ the variation gives us only an upper bound to the true eigenvalue, however, for increasing s the potential and its steepness increases monotonously for any value of ζ as shown in Fig. 2. Thus the kinetic energy and the ground state mass, M_{000}^2 increases with s and therefore for $s > 2$ has to be larger than zero. This can also clearly be seen from the figure where numerical results for $s > 2$ are also shown. For $s \geq 4$ the potential $U_s(\zeta)$ is larger than zero for all values of ζ and therefore M_{000}^2 is positive-definite. Hence the chiral symmetric solution $M_{000}^2 = 0$ is only possible for $s = 2$. For a negative dilaton profile, $\varphi(z) = -\lambda z^s$ the potential is positive semidefinite and no massless pion can be obtained.



FIG. 2: Light-front potential in units $\lambda = 1$ for different powers of s in the dilaton profile (6).

In the following we shall show that the quadratic dilaton profile also follows from the algebraic construction of Hamiltonian operators by dAFF [2]. In fact, the Hamiltonian obtained from unperturbed AdS is identical to the LF Hamiltonian in a free theory of massless quarks in LF quantization [1]. Remarkably such a Hamiltonian can also be derived from a one-dimensional quantum field theory from the action [2]

$$S = \tfrac{1}{2} \int dt \Big(\dot{Q}^2 - \frac{g}{Q^2} \Big), \tag{9}$$

where g is a dimensionless number and $\dot{Q} = \partial_t Q$. The generator of evolution in the variable t, the Hamiltonian, is obtained from (9) by the Noether theorem

$$H_t = \tfrac{1}{2} \left(\dot{Q}^2 + \frac{g}{Q^2} \right). \tag{10}$$

Using the equal-time quantization condition $[Q(t), \dot{Q}(t)] = i$ at $t = 0$, one can express the Hamiltonian H_t in a state-space of functions of one variable, $\psi(x) \in L_2(R_1)$ from the operator representation in the Schrödingier picture $Q \to x$, $\dot{Q} \to -i\frac{d}{dx}$

$$H_t = \tfrac{1}{2}\left(-\frac{d^2}{dx^2} + \frac{g}{x^2}\right). \tag{11}$$

This expression is identical to the free LF Hamiltonian, provided that the variable x is identified with the LF invariant variable ζ, $x = \zeta/\sqrt{2}$, and the Casimir operator of the representation g is related to the LF orbital angular momentum L: $g = L^2 - 1/4$.

The action (9) is invariant under conformal transformations in the variable t, and there are in addition to the Hamiltonian H_t two more invariants of motion for this field theory, namely the dilation operator D and K, corresponding to the special conformal transformations in t. In addition to H_t (10), the operators D and K can be expressed in the field operators $Q(t)$ applying the Noether theorem to the Lagrangian in (9):

$$\begin{aligned} D &= t\,H_t - \tfrac{1}{4}\left(Q\,\dot{Q} + \dot{Q}\,Q\right), \\ K &= t^2\,H_t - \tfrac{1}{2}t\,\left(Q\,\dot{Q} + \dot{Q}\,Q\right) + \tfrac{1}{2}Q^2. \end{aligned} \tag{12}$$

Specifically, if one introduces the the new variable τ defined through $d\tau = dt/(u + v\,t + w\,t^2)$ and the rescaled fields $q(\tau) = Q(t)/(u + v\,t + w\,t^2)^{1/2}$, it then follows that the operator $G = u\,H_t + v\,D + w\,K$ generates evolution in τ [2]

$$G|\psi(\tau)\rangle = i\frac{\partial}{\partial\tau}|\psi(\tau)\rangle. \tag{13}$$

The Hamiltonian corresponding to the operator G is [2]

$$H_\tau = \tfrac{1}{2}\left(\dot{q}^2 + \frac{g}{q^2} + \frac{4\,uw - v^2}{4}q^2\right), \tag{14}$$

where $\dot{q} = \partial_\tau q$.

Noticing that the equal-time quantization condition for Q and $\dot{Q}$ imply the commutation relation $[q(\tau), \dot{q}(\tau)] = i$, we can again represent H_τ in the Schrödinger picture

$$H_\tau = \tfrac{1}{2}\left(-\frac{d^2}{dx^2} + \frac{g}{x^2} + \frac{4\,uw - v^2}{4}x^2\right). \tag{15}$$

For $g \geq -1/4$ and $4\,uw - v^2 > 0$ the resulting operator has a discrete spectrum. Such a Hamiltonian defines in a picture of composite hadrons a bound-state equation. The Hamiltonian H_τ is uniquely determined by the requirement of localization [2], and thus H_τ can be used to study the evolution of the bound-state system in terms of the variable τ. For $x = \zeta/\sqrt{2}$ and $4\,uw - v^2 = 8\lambda^2$ its realization in the space of functions with one variable has exactly the form of the invariant LF Hamiltonian (2) obtained from the AdS metric modified by a dilaton term with a quadratic profile.

The construction of new Hamiltonians from the generators of the conformal group has been used by dAFF to construct algebraically the spectra and the eigenfunctions of these operators. The conformal group in one dimension is locally isomorphic to the group of pseudo-rotations $O(2,1)$. The compact operator $R = \frac{1}{2}(K + H_t)$ generates rotations in the Euclidean 1-2 plane, whereas the non-compact operators $L_1 = \frac{1}{2}(K - H_t)$ and $L_2 = D$ generate pseudo-rotations (boosts) in the non-Euclidean 2-3 and 1-3 plane respectively. As in the familiar case of angular momentum, one can introduce raising and lowering operators $L_\pm = L_1 \pm L_2$ and construct the spectrum and the eigenfunctions analogously to the angular momentum operators. Interestingly, this is just the method employed in Ref. [13] to obtain an integrable LF confining Hamiltonian by following Infeld's observation that integrability follows immediately if the equation of motion can be factorized as a product of linear operators [14]. The method can be extended to describe baryons in AdS while preserving the algebraic structure [13, 15].

To summarize, we have shown in this letter that the triple complementary connection of AdS space, its LF holographic dual, and the relation to the algebra of the conformal group in one dimension is characterized by a quadratic confinement potential, and thus a dilaton profile with the power z^s, with the unique power $s = 2$ and prominent phenomenological features. For $s = 2$ the mass of the $J = L = n = 0$ pion is automatically zero in the chiral limit, and the separate dependence on J and L leads to a mass ratio of the ρ and the a_1 mesons which coincides with the result of the Weinberg sum rules. One predicts linear Regge trajectories with the same slope in the relative orbital angular momentum L and the principal quantum number n. In the phenomenological application constant terms in the potential, which are not fixed by the group theoretical arguments, are very important. They are fixed by the holographic duality to LF quantized QCD [8].

It should be noticed that the overall features of confinement, namely the term proportional to ζ^2 in the effective potential can be derived within a purely group theoretical framework of [2]. It is determined by the term Q^2 in the generator K at $t = 0$, which in turn leads to the quadratic x dependence in the new Hamiltonian (15). Our approach has elements in common with those of Ref. [16], where the scale of the confinement potential arises from a boundary condition when solving Gauss' equation. Notably the confining term x^2 in H_τ is due to the part of the Noether current arising from the total derivative generated by the special conformal transformation. In fact, a possible connection between the construction of different Hamiltonians (evolution operators) and confinement has been already devised by dAFF. Whereas the confining force is usually a consequence of a Lagrangian with dimensionful constants, in the dAFF construction the confining potential is introduced by the choice of the operator, H_τ with $4\,u\,v - v^2 > 0$. The resulting Lagrangian, constrained by the conformal invariance of the action (9), has the same form as the AdS Lagrangian with a quadratic dilaton profile. The AdS approach, however, goes beyond the purely group theoretical considerations of dAFF, since features such as the masslessness of the pion and the separate dependence on J and L are a consequence of the potential (3) derived from the duality with AdS for general J and L.

In their discussion of the evolution operator H_τ as a model for confinement, dAFF mention a critical point, namely that "the time evolution is quite different from a stationary one". By this statement they refer to the fact that the variable τ is related to the variable t for the case $uw > 0$, $v = 0$ by

$$\tau = \frac{1}{\sqrt{u\,w}} \arctan\left(\sqrt{\frac{w}{u}}t\right), \tag{16}$$

i.e., τ has only a limited range. However, we can see from (1) and (13) that the LF time x^+ and the variable τ are related by $\tau = x^+/P^+$ in the LF frame $\vec{P}_\perp = 0$, and thus the finite range of invariant LF time τ can be interpreted as a feature of the internal frame-independent LF time difference between the confined constituents in a bound state. For example, in the collision of two mesons it would allow us to compute the LF time difference between the two possible quark-quark collisions. The LF time $x^+ = P^+\tau$ depends kinematically on the total momentum P^+.

The treatment of the chiral limit in the LF holographic approach to strongly coupled QCD is substantially different from the standard approach based on chiral perturbation theory. In the conventional approach [17], spontaneous symmetry breaking by a non-vanishing chiral quark condensate $\langle\bar{\psi}\psi\rangle$ plays the crucial role. In QCD sum rules [18] $\langle\bar{\psi}\psi\rangle$ brings in non-perturbative elements into the perturbatively calculated spectral sum rules. It should be noted, however, that the definition of the condensate, even in lattice QCD necessitates a renormalization procedure for the operator product, and that it is not a directly observable quantity.

In Ref. [19] it has been argued that quantum field theory is by no means a monolithic block, and that its different facets, aimed at different goals, might well differ in purely theoretical concepts. Condensates are certainly intermediate theoretical concepts inside QCD. The only possibility of direct observation would be by their gravitational interaction, and there, the discrepancy with observation is notorious [20]. Another example is the spontaneous symmetry breaking of the $SU(2) \times SU(2)$ gauge symmetry in the Standard Model. Although the standard treatment of renormalized perturbation theory is based on spontaneously broken symmetry [21–24], the Elitzur theorem [25–27] asserts that a lattice gauge theory cannot be spontaneously broken.

In the confining AdS/QCD model discussed in this paper, the vanishing of the pion mass in the chiral limit, a phenomena usually ascribed to spontaneous symmetry breaking of the chiral symmetry, is obtained specifically from the precise cancellation of the LF kinetic energy and LF potential energy. This provides a viable alternative to the conventional description of nonperturbative QCD based on vacuum condensates; the physics normally ascribed to vacuum phenomena is contained within the light-front wavefunctions of the hadrons themselves [28, 29]. This observation eliminates a major conflict of hadron physics with the empirical value for the cosmological constant.

Acknowledgments

We thank Diego Hofman for pointing out the work of de Alfaro, Fubini, and Furlan to us and to Joseph Day, Paul Hoyer, Shamit Kachru, Ami Katz, Matin Mojaza, and Volkhard Müller for discussions. This work was supported by the Department of Energy contract DE–AC02–76SF00515.

---

[*] sjbth@slac.stanford.edu

[†] gdt@asterix.crnet.cr

[‡] h.g.dosch@thphys.uni-heidelberg.de

[1] G. F. de Teramond and S. J. Brodsky, "Light-front holography: a first approximation to QCD," Phys. Rev. Lett. **102**, 081601 (2009) [`arXiv:0809.4899 [hep-ph]`].

[2] V. de Alfaro, S. Fubini and G. Furlan, "Conformal invariance in quantum mechanics," Nuovo Cim. A **34**, 569 (1976).

[3] A. Karch, E. Katz, D. T. Son and M. A. Stephanov, "Linear confinement and AdS/QCD," Phys. Rev. D **74**, 015005 (2006) [`arXiv:hep-ph/0602229`].

[4] G. F. de Teramond and S. J. Brodsky, "Light-front holography and gauge/gravity duality: the light meson and baryon spectra," Nucl. Phys. Proc. Suppl. **199**, 89 (2010) [`arXiv:0909.3900 [hep-ph]`].

[5] H. C. Pauli, "On confinement in a light cone Hamiltonian for QCD," Eur. Phys. J. C **7**, 289 (1999) [`arXiv:hep-th/9809005`].

[6] S. J. Brodsky, H. C. Pauli and S. S. Pinsky, "Quantum chromodynamics and other field theories on the light cone," Phys. Rept. **301**, 299 (1998) [`arXiv:hep-ph/9705477`].

[7] S. J. Brodsky and G. F. de Teramond, "Hadronic spectra and light-front wavefunctions in holographic QCD," Phys. Rev. Lett. **96**, 201601 (2006) [`arXiv:hep-ph/0602252`].

[8] G. F. de Teramond, H. G. Dosch and S. J. Brodsky, "Kinematical and dynamical aspects of higher-spin bound-state equations in holographic QCD," `arXiv:1301.1651 [hep-ph]`.

[9] G. F. de Teramond and S. J. Brodsky, "Gauge/Gravity duality and hadron physics at the light-front," AIP Conf. Proc. **1296**, 128 (2010) [`arXiv:1006.2431 [hep-ph]`].

[10] P. Breitenlohner and D. Z. Freedman, "Stability in gauged extended supergravity," Annals Phys. **144**, 249 (1982).

[11] S. Weinberg, "Precise relations between the spectra of vector and axial vector mesons," Phys. Rev. Lett. **18**, 507 (1967).

[12] J. W. Strutt, Baron Rayleigh, *The Theory of Sound*, Vol. 1, Dover Reprint New York, 1945, Sec. 88.

[13] S. J. Brodsky and G. F. de Teramond, "AdS/CFT and light-front QCD," World Scientific Subnuclear Series, **45**, 139 (2007) [`arXiv:0802.0514 [hep-ph]`]; G. F. de Teramond, S. J. Brodsky, "Light-front quantization approach to the gauge-gravity correspondence and hadron spectroscopy," AIP Conf. Proc. **1257**, 59 (2010) [`arXiv:1001.5193 [hep-ph]`].

[14] L. Infeld, "On a new treatment of some eigenvalue problems", Phys. Rev. **59**, 737 (1941).

[15] R. P. Martinez y Moreno and A. L. Salas Brito, "Conformal invariance in a Dirac oscillator," J. Math. Phys. **33**.

[16] P. Hoyer, "Relativistic bound states at Born level," `arXiv:1210.8073 [hep-ph]`.

[17] J. Goldstone, A. Salam and S. Weinberg, "Broken symmetries," Phys. Rev. **127**, 965 (1962).

[18] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, "QCD and resonance physics. Sum rules," Nucl. Phys. B **147**, 385 (1979).

[19] H. G. Dosch and V. F. Müller, "The facets of relativistic quantum field theory," Eur. Phys. J. H **35**, 331 (2011).

[20] N. Straumann, "The History of the cosmological constant problem," `arXiv:gr-qc/0208027`.

[21] P. W. Higgs, "Broken symmetries, massless particles and gauge fields," Phys. Lett. **12**, 132 (1964).

[22] F. Englert and R. Brout, "Broken symmetry and the mass of gauge vector mesons," Phys. Rev. Lett. **13**, 321 (1964).

[23] G. S. Guralnik, C. R. Hagen and T. W. B. Kibble, "Global conservation laws and massless particles," Phys. Rev. Lett. **13**, 585 (1964).

[24] G. 't Hooft and M. J. G. Veltman, "Regularization and renormalization of gauge fields," Nucl. Phys. B **44**, 189 (1972).

[25] S. Elitzur, "Impossibility of spontaneously breaking local symmetries," Phys. Rev. D **12**, 3978 (1975).

[26] G. F. De Angelis, D. de Falco and F. Guerra, "A Note on the abelian Higgs-Kibble model on a lattice: Absence of spontaneous magnetization," Phys. Rev. D **17**, 1624 (1978).

[27] G. Scharf. "Quantum Gauge Theory – a True Ghost Story," Wiley, New York, 2001.

[28] S. J. Brodsky and R. Shrock "Condensates in quantum chromodynamics and the cosmological constant," Proc. Nat. Acad. Sci. **108**, 45 (2011) [`arXiv:0905.1151 [hep-th]`].

[29] S. J. Brodsky, C. D. Roberts, R. Shrock , P. Tandy, "New perspectives on the quark condensate," Phys. Rev. **C82**, 022201 (2010) [`arXiv:1005.4610 [nucl-th]`].